

Supplementary Table S3. *C. jejuni* counts (\log_{10} MPN) of seeder and sentinel broiler chickens at 2, 3, 4, 8, 11, 16 and 18 days after inoculation from cloacal swabs.

¹ 2 d post inoculation		² 3 d post inoculation		⁴ 4 d post inoculation		⁸ 8 d post inoculation		¹¹ 11 d post inoculation		¹⁶ 16 d post inoculation		¹⁸ 18 d post inoculation	
control	CE-culture	control	CE-culture	control	CE-culture	control	CE-culture	control	CE-culture	control	CE-culture	control	CE-culture
0.36	2.36	nd	nd	nd	1.36	4.36	4.36	5.36	3.36	6.36	4.36	7.36	3.36
3.36	1.36	nd	nd	nd	nd	5.36	3.36	5.36	3.36	6.36	6.36	6.36	6.36
2.36	1.36	nd	nd	nd	nd	5.36	3.36	6.36	4.36	4.36	3.36	6.36	4.36
0.36	2.36	nd	nd	2.36	nd	5.36	4.36	7.36	5.36	6.36	5.36	8.36	6.36
1.36	2.36	nd	nd	nd	1.36	4.36	3.36	5.36	3.36	5.36	2.36	5.36	4.36
1.36	1.36	nd	1.36	nd	3.36	5.36	4.36	5.36	5.36	5.36	6.36	5.36	4.36
0.36	3.36	nd	nd	nd	nd	5.36	2.36	5.36	5.36	5.36	5.36	5.36	5.36
3.36	2.36	nd	2.36	nd	3.36	4.36	3.36	4.36	5.36	5.36	4.36	5.36	5.36
1.36	1.36	nd	nd	nd	nd	5.36	1.36	5.36	4.36	5.36	5.36	5.36	5.36
3.36	0.36	nd	nd	nd	0.36	7.36	2.36	5.36	3.36	6.36	4.36	6.36	4.36
3.36	3.36	nd	nd	nd	1.36	5.36	4.36	5.36	4.36	5.36	6.36	4.36	5.36
0.36	0.36	nd	nd	nd	nd	5.36	4.36	6.36	5.36	6.36	4.36	5.36	4.36
1.36	2.36	nd	nd	nd	nd	5.36	3.36	6.36	3.36	5.36	5.36	6.36	4.36
2.36	1.36	nd	nd	nd	1.36	6.36	4.36	8.36	4.36	7.36	4.36	6.36	5.36
2.36	2.36	nd	nd	nd	nd	5.36	2.36	6.36	3.36	6.36	4.36	6.36	5.36
2.36	3.36	nd	3.36	nd	3.36	5.36	3.36	6.36	4.36	6.36	5.36	5.36	5.36
3.36	3.36	nd	nd	nd	2.36	5.36	4.36	6.36	3.36	7.36	4.36	7.36	4.36
0.36	2.36	0.36	nd	1.36	1.36	5.36	4.36	5.36	5.36	5.36	4.36	5.36	4.36
		nd	nd	nd	nd	5.36	2.36	8.36	3.36	4.36	5.36	6.36	4.36
		nd	nd	nd	nd	5.36	3.36	6.36	5.36	5.36	4.36	6.36	5.36
		nd	nd	nd	nd	4.36	3.36	4.36	3.36	6.36	4.36	6.36	3.36
		nd	nd	nd	1.36	4.36	4.36	5.36	3.36	5.36	2.36	4.36	4.36
		nd	nd	nd	0.36	3.36	4.36	5.36	3.36	5.36	4.36	5.36	5.36
		nd	nd	nd	1.36	6.36	3.36	6.36	4.36	6.36	4.36	6.36	3.36
		nd	nd	nd	nd	5.36	4.36	7.36	4.36	6.36	4.36	5.36	3.36
		nd	2.36	nd	3.36	5.36	5.36	4.36	5.36	6.36	5.36	6.36	5.36
		4.36	3.36	nd	3.36	4.36	2.36	5.36	3.36	4.36	4.36	4.36	2.36
		nd	nd	nd	1.36	5.36	3.36	5.36	4.36	6.36	4.36	6.36	5.36
		nd	0.36	nd	3.36	4.36	5.36	5.36	4.36	6.36	4.36	6.36	4.36
		nd	nd	nd	nd	5.36	4.36	4.36	3.36	6.36	4.36	6.36	4.36
		nd	3.36	nd	2.36	4.36	3.36	5.36	5.36	6.36	5.36	6.36	4.36
		nd	nd	nd	0.36	4.36	3.36	5.36	3.36	6.36	5.36	5.36	6.36
		nd	nd	nd	3.36	5.36	2.36	5.36	3.36	6.36	4.36	6.36	4.36
		nd	2.36	1.36	3.36	5.36	4.36	6.36	4.36	6.36	5.36	5.36	5.36
		nd	nd	nd	nd	4.36	3.36	4.36	4.36	6.36	3.36	5.36	4.36
		nd	nd	nd	nd	6.36	7.36	5.36	4.36	6.36	5.36	7.36	4.36
median \log_{10} MPN/cloacal swab		1.86	2.36	2.36	2.36	1.36	1.72	5.36	3.36	5.36	4.36	6.36	4.36
Log reduction		-	-	-	-	-	-	2	1	1	2	2	2

¹median values in \log_{10} MPN/g from seeders n = 18; ²median values in \log_{10} MPN/g from sentinels n = 36; "nd" indicates that *C. jejuni* was "not detectable".