

Supplementary material.

Ecology and distribution of red king crab larvae in the Barents Sea: a review

Vladimir G. Dvoretsky* and Alexander G. Dvoretsky

Water

Table S1. Mean abundance (individuals m⁻²) of red king crab larvae in the plankton in Ura Bay in 1996–1999.

Year	Number of samples	Total number of larvae. ind.	March	April	May	June	Mean annual
1996	29	586	36.4	57.4	43.0	7.6	40.4
1997	34	482	9.8	18.0	34.4	-	28.4
1998	38	193	3.4	13.6	12.6	3.4	10.2
1999	201	24	-	2.6	15.4	10.8	12.4

'-' – not found.

Table S2. Total abundance (individuals m⁻²) of different zoea stages (ZI–ZIV) of in the plankton in Ura Bay in 1996–1999.

Year	March				April			
	ZI	ZII	ZIII	ZIV	ZI	ZII	ZIII	ZIV
1996	16.8	13.2	0.8	-	16.2	33.8	2.6	-
1997	6.4	2.8	-	-	6.4	10.0	-	-
1998	3.4	-	-	-	5.4	6.0	-	-
1999	-	-	-	-	8.0	-	-	-
May					June			
1996	10.2	32.6	7.4	1.4	-	-	1.6	4.8
1997	3.2	23.0	5.2	1.0	-	-	-	-
1998	1.4	7.4	2.6	0.4	-	-	-	3.4
1999	1.2	5.6	5.4	0.8	-	-	1.0	8.0

'-' – not found.

Source.

Matyushkin, V.B.; Ushakova, M.F. Larvae of red king crab in the fjords of Western Murman In *The red king crab in the Barents Sea*. Berenboim, B.I., Ed.; PINRO Press: Murmansk, Russia, 2003, pp. 133–140. (In Russian)