

Supplementary Material

Water

# Marine hazard assessment of soluble and nanostructured forms of the booster biocide DCOIT in tropical waters

Fernando Perina <sup>1,2\*</sup>, Cristiane Ottoni <sup>2,3</sup>, Juliana Santos <sup>2</sup>, Vithória Santos <sup>1,4</sup>, Mariana Silva <sup>2</sup>, Bruno Campos <sup>2</sup>, Myana Fontes <sup>2</sup>, Debora Santana <sup>2</sup>, Frederico Maia <sup>5</sup>, Denis Abessa <sup>2</sup> and Roberto Martins <sup>1</sup>

- <sup>1</sup> CESAM-Centre for Environmental and Marine Studies and Department of Biology, University of Aveiro, 3810-193 Aveiro, Portugal.
  - <sup>2</sup> Biosciences Institute, São Paulo State University (UNESP), 11330-900 São Vicente/SP, Brazil.
  - <sup>3</sup> LEAF-Linking Landscape, Environment, Agriculture and Food Research Center, Associated Laboratory TERRA, Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017 Lisboa, Portugal.
  - <sup>4</sup> Institute of Advanced Sea Studies (IEAMAR), São Paulo State University (UNESP), São Vicente/SP, Brazil.
  - <sup>5</sup> Smallmatek - Small Materials and Technologies, Lda, 3810-075 Aveiro, Portugal.
- \* Correspondence: [perinafc@ua.pt](mailto:perinafc@ua.pt)


*\*Corresponding author*

Fernando Cesar Perina

*E-mail address:* [perinafc@ua.pt](mailto:perinafc@ua.pt)

**S1. Standard curve of calibration (absorbance x cell density) - *Chlorella minutissima***

cell/mL	Abs (680 nm)
2500	0.008
5000	0.015
10000	0.025
20000	0.05
40000	0.125
80000	0.192
160000	0.449



**Figure S1.** Absorbance at 680 nm of *Chlorella minutissima* culture as a function of cell concentration.